


OVARIAN CANCER

The Silent Killer - It Whispers, So Listen

Ovarian cancer is expected to be diagnosed in over 295,000 women throughout the world in 2020 with more than 185,000 women dying from this deadly disease⁵ having a 5-year survival rate of less than 50%. Ovarian cancer is often diagnosed in later stages due to its symptoms being misdiagnosed as a more benign condition such as Irritable Bowel Syndrome or Overactive Bladder. Treatment typically requires surgery and chemotherapy.

1 OUT OF 78 WOMEN WILL BE DIAGNOSED WITH OVARIAN CANCER IN THEIR LIFETIME²



Every **24** minutes a woman in the US is diagnosed with ovarian cancer¹



A Pap test is not used to detect ovarian cancer²



www.OverrunOvarianCancer.com

Ovarian cancer can occur in women **At any age**⁴

OVARIAN CANCER IN THE USA

ANNUAL NEW CASES¹ | ANNUAL DEATHS¹
21,404  **14,204**

Ovarian cancer causes more deaths each year in the USA than any other gynecologic cancer⁶

TYPICAL SYMPTOMS OF OVARIAN CANCER³



Bloating



Pelvic or Abdominal Pain



Urinary Urgency or Frequency



Difficulty Eating or Feeling Full Quickly

Risk factors for ovarian cancer:

- Family history
- BRCA1 or BRCA2 gene mutations
- Women who never had children
- Women of Eastern European or Ashkenazi Jewish heritage³



Diagnostic tests for ovarian cancer:

- Transvaginal Ultrasound (TVUS)
- CA-125 blood test
- Pelvic exam

Ovarian cancer research is ongoing, however, currently there are no reliable screening tests².



OVARIAN CANCER RESOURCES

- (1) www.statecancerprofiles.cancer.gov (6) www.cdc.gov/cancer/ovarian/
(2) www.cancer.org
(3) www.sloca.org
(4) www.mayoclinic.org
(5) www.ncbi.nlm.nih.gov/pmc
• www.ocrfa.org/
• OVERRUNovariancancer.com
• American Cancer Society
Helpline 1-800-227-2345



www.HopeLightProject.com